

Light meson spectroscopy with the KLOE experiment.

The KLOE collaboration presented by B. Di Micco ^a,

^a*Dipartimento di Fisica dell'Università "Roma Tre" e Sezione INFN, Roma, Italy.*

Abstract

In this paper we describe the status of the analyses in progress on light meson spectroscopy in the KLOE experiment. We present the analyses of ϕ decays into $f_0(980)\gamma$ and $a_0(980)\gamma$, the Dalitz plot analysis of the $\eta \rightarrow \pi^+\pi^-\pi^0$ decay, the branching ratio measurement of $\eta \rightarrow \pi^0\gamma\gamma$, the upper limits on $Br(\eta \rightarrow 3\gamma)$ and $Br(\eta \rightarrow \pi^+\pi^-)$, the measurement of the ratio $Br(\phi \rightarrow \eta'\gamma)/Br(\phi \rightarrow \eta\gamma)$ and ϕ leptonic width measurements.

1 Introduction

¹ A.Aloisio, F.Ambrosino, A.Antonelli, M.Antonelli, C.Bacci, M.Barva, G.Bencivenni, S.Bertolucci, C.Bini, C.Bloise, V.Bocci, F.Bossi, P.Branchini, S.A.Bulychjov, R.Caloi, P.Campana, G.Capon, T.Capussela, G.Carboni, F.Ceradini, F.Cervelli, F.Cevenini, G.Chiefari, P.Ciambrone, S.Conetti, E.De Lucia, A.De Santis, P.De Simone, G.De Zorzi, S.Dell'Agnello, A.Denig, A.Di Domenico, C.Di Donato, S.Di Falco, B.Di Micco, A.Doria, M.Dreucci, O.Erriquez, A.Farilla, G.Felici, A.Ferrari, M.L.Ferrer, G.Finocchiaro, C.Forti, P.Franzini, C.Gatti, P.Gauzzi, S.Giovannella, E.Gorini, E.Graziani, M.Incagli, W.Kluge, V.Kulikov, F.Lacava, G.Lanfranchi, J.Lee-Franzini, D.Leone, F.Lu, M.Martemianov, M.Martini, M.Matsyuk, W.Mei, L.Merola, R.Messi, S.Miscetti, M.Moulson, S.Müller, F.Murtas, M.Napolitano, F.Nguyen, M.Palutan, E.Pasqualucci, L.Passalacqua, A.Passerì, V.Patera, F.Perfetto, E.Petrolo, L.Pontecorvo, M.Primavera, P.Santangelo, E.Santovetti, G.Saracino, R.D.Schamberger, B.Sciaccia, A.Sciubba, F.Scuri, I.Sfiligoi, A.Sibidanov, T.Spadaro, E.Spiriti, M.Testa, L.Tortora, P.Valente, B.Valeriani, G.Venanzoni, S.Veneziano, A.Ventura, R.Versaci, I.Villella, G.Xu.

² Corresponding author: B. Di Micco, Università "Roma Tre", Via della Vasca Navale, 84, I-00146, Roma, Italy, e-mail dimicco@fis.uniroma3.it

The KLOE detector [1], operates at the Frascati e^+e^- collider DAΦNE [2], which runs at a CM energy W equal to the ϕ -meson mass, $W \sim 1019.5$ MeV. The analyses presented here are based on data collected in the years 2001 and 2002 for an integrated luminosity of $\sim 450 \text{ pb}^{-1}$ corresponding to 1.5 billions of ϕ and 20 millions of η mesons [$Br(\phi \rightarrow \eta\gamma) \sim 1.3\%$ [3]]. This means that KLOE can also study η physics in a clean environment with high statistic.

2 Search for $\phi \rightarrow f_0\gamma$ in $\pi^+\pi^-\gamma$ events.

The ϕ radiative decays to scalar mesons, $\phi \rightarrow S\gamma$, give significant insight in the assessment of the nature of lower mass scalar mesons [4]. With the KLOE experiment the decays $\phi \rightarrow f_0(980)\gamma$ and $\phi \rightarrow a_0(980)\gamma$ are searched for in $\pi^0\pi^0\gamma$ and $\eta\pi^0\gamma$ [5,6] final states respectively. Moreover the $f_0(980)$ signal is also searched for $\pi^+\pi^-\gamma$ events with a photon at large angle. The search for this signal is characterized by the presence of a huge irreducible background due to the initial state radiation (ISR), to $e^+e^- \rightarrow \pi^+\pi^-\gamma$ (FSR) and $\phi \rightarrow \rho^\pm(\rightarrow \pi^\pm\gamma)\pi^\pm$. The f_0 events are searched for in the large photon angle region $45^\circ < \theta < 135^\circ$ to reduce ISR background. The f_0 signal appears as a bump in the $\pi^+\pi^-$ invariant mass $M_{\pi\pi}$ spectrum around 980 MeV. Fig.1 (top) shows the spectrum obtained at $\sqrt{s} = M_\phi$.

An overall fit to the spectrum has been done by applying the following formula:

$$\frac{dN}{dM_{\pi\pi}} = \left[\left(\frac{d\sigma}{dM_{\pi\pi}} \right)_{ISR} + \left(\frac{d\sigma}{dM_{\pi\pi}} \right)_{FSR+f_0} + \left(\frac{d\sigma}{dM_{\pi\pi}} \right)_{\rho\pi} \right] \times L \times \epsilon(M_{\pi\pi})$$

with L the integrated luminosity and $\epsilon(M_{\pi\pi})$ the selection efficiency as a function of $M_{\pi\pi}$. The f_0 amplitude is taken from the kaon-loop approach [7]. A forward-backward asymmetry $A = \frac{N^+(\theta > 90^\circ) - N^+(\theta < 90^\circ)}{N^+(\theta > 90^\circ) + N^+(\theta < 90^\circ)}$ is expected, due to the interference between FSR

and ISR[8]. Fig.1 (bottom) shows the asymmetry as a function of $M_{\pi\pi}$ compared to the prediction based on the ISR-FSR interference alone. A significant deviation from the prediction is observed in the high mass region clearly due to the f_0 contribution.




Fig. 1. (top) $M_{\pi\pi}$ spectrum of $\pi^+\pi^-\gamma$. The upper (lower) curves are the result of the fit and the estimated background due to ISR, FSR and $\rho\pi$. (bottom) Forward-Backward asymmetry A as a function of $M_{\pi\pi}$. The curve and the black points are the Montecarlo expectations based on the interference between FSR and ISR only. The experimental data are reported as triangles.

3 Dynamics of $\eta \rightarrow \pi^+ \pi^- \pi^0$

The dynamics of the $\eta \rightarrow \pi^+ \pi^- \pi^0$ decay has been studied with a Dalitz plot analysis. The conventional variables X and Y are defined as: $X = \sqrt{3} \frac{T_+ - T_-}{Q_\eta}$, $Y = \frac{3T_0}{Q_\eta} - 1$, where $Q_\eta = m_\eta - 2m_{\pi^+} - m_{\pi^0}$ and T_+ , T_- and T_0 are the kinetic energies of the particles. The measured distribu-

tion has been fitted as: $|A(X, Y)|^2 \simeq (1 + aY + bY^2 + cX + dX^2 + eXY + \dots)$. C-parity conservation prevents odd powers in X being present in the expansion: thus parameters c and e should be zero as confirmed by our fit. The results of the fit are shown in table 3 Efficiency is $\sim 36\%$ over

Table 1

Fitted parameters $P(\chi^2) = 52\%$ of $\eta \rightarrow 3\pi$ Dalitz plot.

a	b	c
$-1.075 \pm .008$	$.118 \pm .009$	$-.0005 \pm .004$
d	e	f
$.049 \pm .008$	$-.004 \pm .01$	$.13 \pm .02$

the whole Dalitz plot. The evaluation of systematic effects is under completion.

4 Rare and forbidden η decays

$$(\eta \rightarrow \pi^0 \gamma \gamma, \eta \rightarrow \pi^+ \pi^-, \eta \rightarrow \gamma \gamma \gamma)$$

The $\eta \rightarrow \pi^0 \gamma \gamma$ decay is interesting to test the Chiral Perturbation Theory prediction for the branching ratio and $m_{\gamma\gamma}$ spectrum[9,10]. The most accurate measurement for the branching ratio[11] is, infact, far from any theoretical prediction for this decay based on ChPT. Recently a new measurement has been performed [12] giving a much lower value than the previous one, with a larger error. All previous experiments were done at hadron machines, using mainly $\pi^- p \rightarrow \eta n$, and are largely dominated by $\pi^0 \pi^0$ background and geometrical acceptance. KLOE performs a measurement of competitive precision in a cleaner environment.

Furthermore, it has different background topologies and experimental systematics. The signal is searched for by looking for a $\pi^0 \gamma \gamma \gamma$ topology, where the further γ comes from $\phi \rightarrow \eta \gamma$. Five prompt clusters are required and an overall kinematic fit requiring π^0 mass is performed. The clusters energy must be > 30 MeV and azimuthal angle $> 21^\circ$ to reject fake clusters coming from machine background. The dominant background channel is $\eta \rightarrow 3\pi^0$ that had been reduced with several topological cut. With this selection we obtain an efficiency of 5.7% . To give an idea of the sensitivity, In fig. 2 we compare the $M(4\gamma)$ data spectrum with MC based predictions of signal and background in two hypotheses for the size of signal: one based on PDG value[11] and one based on the recent CB result[12]. It is evident that our data are incompatible with [11] and are marginally in agreement with [12]. The background simulation and the efficiency for the signal is still under study.

$\eta \rightarrow 3\gamma$ decay is C violating. It is a sensitive test of C violation in the strong and electromagnetic interactions. For the details of this analysis see ref.[13]. The KLOE result for the branching ratio is: $Br(\eta \rightarrow \gamma \gamma \gamma) \leq 1.6 \times 10^{-5}$ @90 % C.L. This limit is the best experimental limit for this decay. The expected branching ratio from the Standard Model is $\leq 10^{-12}$ [14], so any discovery of a larger decay rate would be a clear signal of Standard Model deviation.




Fig. 2. $M(4\gamma)$, the spectra expected from the GAMS[11] and Crystall Ball[12] measurement are shown. In the lower plot we show also the expected spectrum for a $Br \sim 1/2$ of C.B. result.

$\eta \rightarrow \pi^+\pi^-$ decay is P and CP violating. This decay is allowed as a weak direct CP violating decay with a very low branching ratio: $BR(\eta \rightarrow \pi^+\pi^-) \sim 10^{-27}$ [15]. Therefore the detection of this decay at an accessible level would be a signal of P and CP violation not explainable in the Standard Model framework. The latest published [16] direct search of this decay has given the following 90% C.L. upper limit: $BR(\eta \rightarrow \pi^+\pi^-) < 3.3 \times 10^{-4}$. In KLOE the signal is searched for the $M(\eta)$ region of the $\pi^+\pi^-$ invariant mass spectrum of $\pi^+\pi^-\gamma$ events selected according to the

$f_0(980) \rightarrow \pi^+\pi^-$ analysis described before (see fig. 1). The signal efficiency is: $\epsilon_s = 16.6\%$. The expected signal has a Gaussian shape with a mass resolution of 1.33 MeV. No signal is observed. The background is determined by fitting the theoretical model for $\pi^+\pi^-\gamma$ sample to the full spectrum. In order to determine an upper limit, we have added to this background a Gaussian function representing the signal multiplied by a constant N_s . We obtain: $N_s = -8 \pm 24$. The 90% confidence level upper limit on the number of events is obtained using the tables in [17]: $N_s < 33$. The branching ratio is $BR(\eta \rightarrow \pi^+\pi^-) = \frac{N_s}{\epsilon_s N_\eta}$ with N_η the number of η in the sample (1.55×10^7). The 90% C.L. upper limit is: $BR(\eta \rightarrow \pi^+\pi^-) < 1.3 \times 10^{-5}$. It improves by a factor ~ 30 the current PDG limit.

5 $\eta - \eta'$ mixing

Here we present the $R = \frac{\Gamma(\phi \rightarrow \eta'\gamma)}{\Gamma(\phi \rightarrow \eta\gamma)}$ measurement. The η' is identified via the decays: $\phi \rightarrow \eta'\gamma$; $\eta' \rightarrow \pi^+\pi^-\eta$; $\eta \rightarrow \pi^0\pi^0\pi^0$ and the decays $\phi \rightarrow \eta'\gamma$; $\eta' \rightarrow \pi^0\pi^0\eta$; $\eta \rightarrow \pi^+\pi^-\pi^0$. The final state is thus characterized by two charged pions and seven photons, and has no physics background with the same topology in KLOE. After background subtraction we observe $3405 \pm 61 \pm 31$ $\phi \rightarrow \eta'\gamma$ events. We normalize to the number of observed $\eta \rightarrow \pi^0\pi^0\pi^0$ decays in the same runs to obtain a preliminary measurement of the ratio of BR's: $R = (4.9 \pm 0.1_{stat} \pm 0.2_{syst}) \times 10^{-3}$.

This result compares favourably with our previous estimate [18] (which already dominates the world average [3]) but with considerably improved accuracy.

6 A new measurement of the ϕ leptonic width.

KLOE has performed a new measurement of the ϕ leptonic widths Γ_l with $l = e, \mu$ [19], using the two data samples taken below ($\sqrt{s}=1017$ MeV) and above ($\sqrt{s}=1022$ MeV) the ϕ peak together with the data taken at the ϕ peak. The dependences on \sqrt{s} of the forward-backward asymmetry of Bhabha events A_{FB} and of the $e^+e^- \rightarrow \mu^+\mu^-$ cross-section $\sigma(\mu\mu)$ around the ϕ peak are sensitive to the value of Γ_{ee} and $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$ respectively. We measure the ϕ mass M_ϕ , the forward-backward asymmetry at $W = M_\phi$ A_{FB}^0 , and finally Γ_{ee} . The result for Γ_{ee} is: $\Gamma_{ee} = 1.32 \pm 0.05_{stat} \pm 0.03_{syst}$ keV. The result for $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$ is: $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}} = 1.320 \pm 0.018_{stat} \pm 0.017_{syst}$ keV. The two results are in good agreement consistently with lepton universality. Combining them we get: $\Gamma_l = 1.320 \pm 0.017_{stat} \pm 0.015_{syst}$ keV with a total uncertainty below 2 %. We point out that the value of Γ_{ee} is necessary for ϕ decay branching ratio measurements, and play also a role in the evaluation of the hadronic contribution to vacuum polarization [20].

References

- [1] KLOE Coll., M. Adinolfi *et al.*, Nucl. Instrum. Meth. **A488**, 51 (2002).
- [2] S. Guiducci, in: Proc. 2001 Particle Accelerator Conf. (ed. P. Lucas, S. Webber, Chicago, 2001).
- [3] K. Hagiwara *et al.*, Phys. Rev. **D66** (2002).
- [4] N.N.Achasov, V.N.Ivanchenko, Nucl.Phys. **B315**, 465 (1989); F.Close, A.Kirk, Phys. Lett. **B515**, 13 (2001); L.Maiani, F.Piccinini, A.D. Polosa, V.Riquer, hep-ph/0407017.
- [5] KLOE Coll., A. Aloisio *et al.*, Phys. Lett. **B536**, 209 (2002).
- [6] KLOE Coll., A. Aloisio *et al.*, Phys. Lett. **B537**, 21 (2002).
- [7] N.N.Achasov, V.V.Gubin, Phys. Rev. **D57**, 1987 (1998).
- [8] S.Binner, J.H.Kuhn, K.Melnikov, Phys. Lett. **B459**, 279 (1999).
- [9] J. Bijnens and J. Gasser, in: Proc. Workshop on Production, Interaction and Decay of the η Meson (ed. J. Bijnens, G. Fäldt, B. M. K. Nefkens, Uppsala, October 2001), **T99**, 34 (Phys. Scripta, Stockholm, 2002).
- [10] E.Oset, J.R.Pelaez, L.Roca, Phys. Rev. **D67**, 073013 (2003).
- [11] Alde *et al.*, Z. Phys. **C25**, 225 (1984).
- [12] N. Knecht *et al.*, Phys. Lett. **B589**, 14 (2004).
- [13] KLOE Coll., A. Aloisio *et al.*, Phys. Lett. **B591**, 49 (2004).

- [14] P. Herczeg, in: Proc. Int. Workshop on Production and Decay of Light Mesons (ed. P. Fleury, Paris) 16 (World Scientific, 1988).
- [15] E. Shabalin,
in: Second DAFNE physics Handbook (ed. Rome, 1995) 445.
- [16] R.R. Akhmetshin *et al.*, Phys.Lett. **B462**, 371 (1999).
- [17] G.J. Feldman, R.D. Cousins, Phys. Rev. **D57**, 3873 (1998).
- [18] KLOE Coll., A. Aloisio *et al.* Phys. Lett. **B541**, 45 (2002).
- [19] M. Antonelli, M. Dreucci, Leptonic decay widths of the phi-meson, KLOE note n. 193 (<http://www.lnf.infn.it/kloe>);
- [20] S. Eidelman, F. Jegerlehner Z.Phys. C67 (1995) 585.

